

Cite this: *Green Chem.*, 2018, 20, 1218

Received 28th December 2017,
Accepted 15th February 2018

DOI: 10.1039/c7gc03880a

rs.c.li/greenchem

An integrated process for enzymatic catalysis allowing product recovery and enzyme reuse by applying thermoreversible aqueous biphasic systems†

Ana M. Ferreira,  ‡^a Helena Passos,  ‡^a Akiyoshi Okafuji,^b Ana P. M. Tavares,^a Hiroyuki Ohno,  ^b Mara G. Freire  *^a and João A. P. Coutinho  *^a

Herein, thermoreversible aqueous biphasic systems (ABS) composed of ammonium-based zwitterions (ZIs) and polymers are disclosed to act as integrated bioreaction–separation processes. The biocatalytic reaction involving laccase occurs in homogeneous media, after which small changes in temperature induce the formation of two phases and the complete separation of the enzyme from the products in a single-step. These systems also allow the recovery and reuse of the enzyme, along with the ZI-rich phase, contributing towards the development of sustainable biocatalytic processes.

In the past few years, there has been increased interest in the development of sustainable and efficient production processes, in which biocatalysis plays a pivotal role.¹ Enzymes display high activity, specificity and selectivity, minimizing the consumption of raw materials and the production of by-products when compared to their synthetic counterparts, thus contributing towards the development of sustainable catalytic processes.^{1,2} Enzymes and biocatalysis have been used in several sectors, such as in soil bioremediation, nanobiotechnology, and biosensors, and in the textile, pulp and paper, cosmetic, food and pharmaceutical industries.³ Significant developments have been made in the pharmaceutical industry with a decrease of the respective *E*-factor.⁴ Despite their advantages, the major challenges associated with enzyme-catalysed bioprocesses rely on the maintenance of the enzyme structural stability and activity, and on their reusability.⁵ Efforts have been made to overcome these issues, whereas one of the main strategies is the enzyme immobilization where the biocatalyst is adsorbed on, bound to, or encapsulated in a solid matrix.^{6,7}

Albeit this approach allows the reuse of the biocatalyst, immobilization may present some disadvantages, such as conformational changes of the enzymes, loss of activity, and their heterogeneity on the support.^{6,8} Therefore, new approaches to recover biocatalysts from the reaction medium have been proposed, including for instance the use of membranes, centrifugation, liquid–liquid extraction,⁹ and the immobilization of chemically modified enzymes.^{7,10}

Liquid–liquid systems appear as an interesting alternative in biocatalysis since they could allow the integration of the reaction and separation steps. Although liquid–liquid systems have been widely employed in the chemical industry for separation purposes due to their simplicity, low cost, and easy scale-up,¹¹ the commonly employed volatile organic solvents can be harmful to biologically active biomolecules. However, a milder environment to biomolecules can be afforded by water-rich systems, such as aqueous biphasic systems (ABS).¹² ABS are composed of two immiscible aqueous-rich phases and can be formed by a mixture of two water soluble phase-forming components, such as two polymers, a polymer and a salt, or two salts, dissolved in aqueous media.¹³ Polymer-based ABS have shown to be biocompatible to deal with the separation of proteins, enzymes, viruses, cells, cell organelles, and other biological materials.¹⁴ Nevertheless, these systems are limited by a narrow polarity range preventing high extraction efficiencies and selectivities. This drawback may be overcome by ABS composed of ionic liquids (ILs).^{12,15} Compared to the more conventional polymer-based systems, IL-based ABS display a lower viscosity, a faster phase separation, and higher extraction performance and selectivity.¹²

Previous studies on IL-based ABS have focused the determination of the respective phase diagrams and the evaluation of their extraction efficiency for a wide range of compounds.¹² Recently, there has been significant interest in the development of dynamic and reversible IL-based ABS induced by the change of pH,¹⁶ temperature¹⁷ or by the addition of gases,^{17,18} that could ultimately lead to the development of integrated reaction–separation processes. However, only a few studies

^aCICECO – Aveiro Institute of Materials, Chemistry Department, University of Aveiro, 3810-193 Aveiro, Portugal. E-mail: maragfreire@ua.pt, jcoutinho@ua.pt

^bDepartment of Biotechnology and Life Science, Tokyo University of Agriculture and Technology, Koganei, Tokyo, Japan

†Electronic supplementary information (ESI) available: Materials and experimental procedures, binodal weight fraction data and phase diagram representation. See DOI: 10.1039/c7gc03880a

‡These authors contributed equally to this work.

investigated the thermoreversible behaviour of IL-based ABS in separation processes. They have been investigated for the extraction of proteins (by temperature change),¹⁹ and in the production and separation of 5-hydroxymethylfurfural (HMF) from fructose (by pH change).²⁰

In addition to the widely studied IL-based systems, novel ABS composed of water soluble ammonium-based zwitterions (ZIs) and inorganic salts have been recently proposed,²¹ and their application in the selective separation of amino acids has been demonstrated. ZIs were suggested as substitutes of ILs, where the cation and the anion are covalently tethered, thus avoiding possible ion exchange between the ABS coexisting phases.²² However, the salts employed may lead to biocompatibility concerns when dealing with solutes sensitive to ionic strength, like proteins and enzymes.²³ To minimize these issues, salts are here proposed to be replaced with polymers, such as polyethylene glycol (PEG), generally recognized as safe (GRAS) and used as additive in food products and drugs.^{10,24,25} Furthermore, ZI-based reversible ABS have not previously been considered as potential integrated reaction-separation platforms.

Based on the need for finding integrated reaction-separation processes for biocatalysis, in this work we investigated and designed novel ZI-polymer-based ABS with thermal switchable behaviour, at temperatures amenable to deal with proteins. These systems were tested as integrated reaction-separation platforms, where a homogeneous catalytic reaction followed by the enzyme separation from the product, by a change in temperature, is performed in one-step. To this end, laccase (EC 1.10.3.2) was used as the biocatalyst and 2,2'-azinobis(3-ethylbenzthiazoline-6-sulfonate) (ABTS) as the substrate. Laccase was chosen since it has been attracting a great deal of attention in the past few years due to its ability to degrade a wide variety of substrates, being currently considered as a promising biocatalyst in the bio-based industries.^{6,26}

Three water-soluble ammonium-based ZIs – *N,N,N*-tripentyl-3-sulfonyl-1-propaneammonium ($N_{555}C3S$), *N,N,N*-tripropyl-3-sulfonyl-1-propaneammonium ($N_{333}C3S$), and *N,N,N*-trimethyl-3-sulfonyl-1-propaneammonium ($N_{111}C3S$), with different alkyl chain lengths, were synthesized and used in the preparation of ABS. Their molecular structures are depicted in Fig. 1. Their synthesis was previously described,¹³ while details on their purity are given in the ESI.†

The ability of each ZI to form ABS when mixed with PEGs of different molecular weights (1540, 2000, 4000 and 6000 g mol⁻¹) was firstly evaluated, and the respective ternary phase diagrams were determined at 25 °C. Further details on the experimental procedure, as well as on the detailed experimental weight fraction data are given in the ESI (Tables S1–S5†). Both $N_{555}C3S$ and $N_{111}C3S$ ZIs are able to form ABS at 25 °C with all polymers studied; however, $N_{333}C3S$, the ZI of intermediate alkyl chain length, does not undergo phase separation with any of the investigated polymers. A summary of the ZI's ability to form ZI-PEG-based ABS is provided in the ESI (Table S6†). To the best of our knowledge, this irregular trend on the ABS formation with an increase of the alkyl side




Fig. 1 (A) Chemical structures and acronyms of the ZIs used. (B) Chemical structure of PEG with the molecular formula H-(O-CH₂-CH₂)_n-OH.

chain length of one of the phase-forming components was never observed, and is certainly a result of a change in the balance of competing interactions between the compounds present in the investigated ternary systems due to the increase of the ZI alkyl chains.^{27,28}

By plotting all the phase diagrams at 25 °C (*cf.* ESI, Fig. S1†), it is possible to appraise the influence of the length of the aliphatic moiety of the ZI, as well as of the molecular weight of the PEG on the solubility curves. The solubility curves indicate the minimum composition required to form ABS, where mixtures with compositions above the curve are biphasic while those below are monophasic. The larger the biphasic region, the higher the ability of the ZI to induce liquid-liquid phase separation in polymer aqueous solutions. ZIs with shorter alkyl side chains are more prone to form ABS with polymers, while the ability of PEG to induce phase separation increases with its molecular weight, which is in agreement with trends previously observed for ABS composed of polymers/salts²⁹ or polymers/ILs.³⁰ This is a consequence of the increase in the hydrophobicity of PEGs with a higher molecular weight.³⁰

After the previous assessment on the ZI-PEG potential to form ABS at a common temperature (25 °C), these were then appraised at 35 and 45 °C, taking into account the operational temperature range of most enzymes and to avoid their activity loss. The optimum operation temperature of laccase is around 40 °C,³¹ although this temperature can vary with the pH and the laccase source, between 30 and 50 °C.³² Fig. 2 depicts the phase diagrams obtained for ZI-PEG 6000-based ABS at several temperatures. The detailed experimental procedure used and the experimental data obtained for PEG 6000 and remaining polymers at different temperatures are provided in the ESI (Fig. S2†). It should be noted that $N_{555}C3S$ does not form ABS with PEG 6000 at 45 °C. As previously observed for ZI-salt-based ABS,²¹ ZI-polymer-based ABS also present two distinct behaviours with temperature, which depend on the ZI alkyl chain length: (i) an increase of the immiscibility region with an increase in temperature for $N_{111}C3S$ -based ABS; and (ii) a decrease of the biphasic region with a decrease in temperature




Fig. 2 Temperature effect in the phase diagrams of ternary systems composed of ZI + PEG 6000 + H₂O at 25 °C (\blacktriangle), 35 °C (\bullet) and 45 °C (\blacksquare). (A) N₁₁₁C3S- and (B) N₅₅₅C3S-based ABS. There is no formation of ABS comprising PEG 6000 and N₅₅₅C3S at 45 °C.

for the N₅₅₅C3S-based systems. ABS composed of ZIs with smaller alkyl chains (N₁₁₁C3S) display a lower critical solution temperature (LCST)-type behaviour when mixed with polymers, following the same behaviour as that of polymer-salt ABS.³³ Their formation thus seems to be dominated by a salting-out phenomenon, where the ZI acts as the salting-out species.³⁰ On the other hand, the ABS formed with the most hydrophobic ZI investigated – N₅₅₅C3S – present an upper critical solution temperature (UCST)-type behaviour, similar to those of polymer-polymer ABS.³⁴ At higher temperatures, the interactions between the polymer and ZI are more favourable, thus enhancing their mutual solubility and reducing their ability to form ABS.³⁵

The temperature-reversible behaviour of the studied systems is also depicted in Fig. 2. To confirm this behaviour, a ternary mixture at the monophasic region was prepared at 25 °C for the system composed of N₁₁₁C3S, and at 45 °C for the system composed of N₅₅₅C3S. Then, the temperatures of these systems were changed to 45 °C and 25 °C for the systems composed of N₁₁₁C3S and N₅₅₅C3S, respectively, to induce the formation of a biphasic system. Details on these mixture compositions are described below, corresponding to the same mixtures at which biocatalysis was carried out. When the phase separation occurs, a top polymer-rich phase and a bottom ZI-rich phase are formed. This proved the thermoreversible

behaviour of the studied systems which were then applied to develop integrated reaction–separation biocatalytic processes. Herein, the homogeneous biocatalytic oxidation of ABTS by laccase was carried out in the monophasic region, followed by the separation of laccase from the oxidation product by promoting the phase separation by temperature changes (increase or decrease according to the ZI employed in the ABS formation). The oxidation of ABTS occurred through the addition of laccase and ABTS to a monophasic mixture composed of ZI, PEG 6000 and water. The mixture was carefully mixed for 1 min, at 40 °C for ZI N₁₁₁C3S (20 wt% ZI + 20 wt% PEG 6000), and at 25 °C for ZI N₅₅₅C3S (39 wt% ZI + 13 wt% PEG 6000) – experimental details are given in the ESI.† The negligible effect of laccase, ABTS and PBS in the phase diagram behaviour and ABS phase compositions was confirmed by the determination of an additional phase diagram in the presence of all these components at the concentrations used in the separation assays (*cf.* ESI, Fig. S2†).

For the two studied thermoreversible ABS, it was observed that laccase successfully catalysed the oxidation of ABTS (colourless) to a highly stable green-coloured ABTS⁺ radical³⁶ (Fig. 3A). ABTS is a standard substrate used to appraise the laccase activity, for which the yield of the reaction is considered as 100%. After that, a temperature change (from 25 to 40 °C or *vice versa*) was applied, inducing the separation of the system into two phases, while allowing the selective separation of the oxidized substrate and laccase, as shown in Fig. 3. Further experimental details are given in the ESI.†

Fig. 3B shows the extraction efficiencies (EE%) of the studied systems for both laccase and oxidized ABTS, which correspond to the percentage recovery of each compound to a given phase. Remarkably, laccase and the reaction product partition to opposite phases, with extraction efficiencies higher than 81% achieved in a single-step. For the two studied ABS, laccase preferentially partitions to the ZI-rich phase and the reaction product to the PEG-rich phase, allowing thus the selective separation of the enzyme and product. The trend obtained for the oxidized ABTS is in good agreement with its hydrophobic character ($(\log(K_{ow}) = 1.93)$,³⁷ where K_{ow} is defined as the octanol–water partition coefficient), meaning that ABTS tends to enrich in more hydrophobic phases, corresponding to the polymer-rich phase. On the other hand, proteins and enzymes tend to preferentially migrate to more hydrophilic (water-rich) phases, and particularly laccase as demonstrated in the literature,^{38,39} therefore preferentially migrating to the ZI-rich phase. Previous studies on the separation of proteins using ABS formed by ILs and polymers support this behaviour.^{40,41}

A remarkable extraction efficiency of 100% was obtained with the N₁₁₁C3S-based ABS, *i.e.* laccase and the oxidized ABTS are completely separated for opposite phases with no losses or cross-contamination. It should be remarked that complete separation could also be achieved with other ZI-based ABS by a proper tailoring of the hydrophobicity of phases, either by changing the PEG molecular weight, ZI employed, or the mixture composition.




Fig. 3 (A) Oxidation of ABTS using laccase and their selective separation from the reaction product by changes in temperature. (B) Extraction efficiencies of laccase (EE%, orange bars) and extraction efficiencies of the green-coloured ABTS⁺ radical (EE%, green bars) in ABS formed by PEG 6000 and N₅₅₅C3S or N₁₁₁C3S.

In addition to the complete separation of the enzyme and the reaction product, it was further confirmed that laccase maintains its biocatalytic activity after the separation step (experimental details and results are given in the ESI[†]). This maintenance in activity is of crucial relevance when envisaging the development of processes in which the enzyme can be recovered and reused. To prove this concept, the phase enriched in laccase (ZI-rich) was recovered and reused for the creation of a novel ternary system, as a new biocatalytic reaction medium. A summary of the enzyme recovery/reuse process is depicted in Fig. 4A; experimental details are given in the ESI[†]. Remarkably, it is possible to recover laccase and reuse it without losses in the catalytic activity, for at least 5 consecutive cycles (Fig. 4B). This approach not only allows the recovery and reuse of the enzyme, but also allows the recovery and reuse of the ZI-rich phase required to form ABS, contributing towards the development of sustainable biocatalytic processes. If a sustainable method is being proposed, the PEG-

rich phase should also be recovered and reused. An ultrafiltration step was finally applied to the PEG-rich phase, allowing the recovery of the oxidized ABTS and ZI from the polymer-rich solution (experimental details are given in the ESI[†]), showing that it is possible to recover the polymer with negligible losses (*cf.* ESI, Fig. S3 and S4[†]).

Two studies were previously reported using two-phase systems with laccase as a biocatalyst.^{42,43} In the first work, it was shown the occurrence of enzymatic reactions (using laccase from *Polyporus versicolor*) in a two-phase system, yet composed of one organic solvent, such as hexane, toluene, tetrahydrofuran, acetone, among others, comprising the substrate and products, and an aqueous phase which contains the enzyme.⁴² However, the enzymatic reaction occurs in heterogeneous media, *i.e.* at the interface, thus being essentially different from the approach here proposed. In addition to the use of hazardous organic solvents, the authors³⁶ were not able to characterize the compounds obtained in aqueous solution




Fig. 4 (A) Flowchart of the integrated reaction–separation process developed, including the enzyme and ZI-rich phase recyclability. (B) Relative laccase activity in the ZI-rich phase in 5 cycles of oxidative reaction, comprising both the recovery and reuse of the enzyme and ZI-rich phase.

due to difficulties in obtaining them in sufficient amounts. Nicotra *et al.*⁴³ isolated and characterized four C–C and C–O dimers from the steroid hormone 17 β -estradiol, formed by a catalytic reaction using laccase from *Trametes pubescens* in a biphasic system, formed by ethyl acetate and an aqueous buffered salt solution. Also in this approach, biocatalysis occurred at the interface. These heterogeneous reaction media usually require vigorous shaking and stirring to ensure the transfer of substrates and products between the phases, which can cause enzyme inactivation.⁴⁴ This may be responsible for the low yield obtained, *ca.* 27%.⁴³ Andersson and Hahn-Hägerdal⁹ reviewed other studies on bioconversion in conventional (polymer–salt and polymer–polymer) ABS, using different enzymes or living cells, but in all these cases ABS were used as heterogeneous media, *i.e.* with systems involving two-phases with the reaction occurring at the interface. Moreover, in these studies, it was not possible to adequately separate the enzyme and the product for opposite phases. To overcome this problem, the use of ultrafiltration units has been proposed.⁹ The thermo-reversible systems here suggested thus overcome these limitations, allowing a homogeneous biocatalytic reaction, with all the advantages in terms of kinetics and yields, followed by a complete separation of the products and enzymes by a small temperature change.

In summary, temperature-driven reversible ABS composed of ZIs and polymers are efficient and sustainable integrated reaction–separation platforms in biocatalytic processes. The reaction and separation steps can be carried out sequentially by taking advantage of their temperature switchable behaviour. The reaction step occurs in a homogeneous medium, followed by the enzyme and product separation in liquid–liquid systems promoted by small changes in temperature. This approach avoids the need for vigorous stirring to improve mass transfer, as typically carried out in heterogeneous reactions, thus contributing to the maintenance of the enzyme activity. Furthermore, the thermoreversible nature of ZI-PEG-ABS is observed at temperatures amenable for maintaining the enzyme stability and activity (25–45 °C). These systems further allow the complete separation of the enzyme and the products in one-step, which are enriched in opposite phases, followed by the recovery and reuse of both the ZI-rich phase and enzyme. In addition to the example discussed here, reversible ZI-based ABS can be tailored to fit the requirements of other biocatalytic processes comprising value-added products.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was developed in the scope of the project CICECO-Aveiro Institute of Materials (Ref. FCT UID/CTM/50011/2013), financed by national funds through the FCT/MEC

and when applicable co-financed by FEDER under the PT2020 Partnership Agreement. The authors also acknowledge FCT for the PhD grants SFRH/BD/85248/2012 and SFRH/BD/92200/2013 of H. P. and A. M. F., respectively. A. M. F. and H. P. thank the financial support of Professor H. Ohno during their stay in his laboratory for collaboration work. A. P. M. Tavares acknowledges FCT for the Investigator Programme (IF/01634/2015). The authors acknowledge financial support from the European Union Framework Programme for Research and Innovation HORIZON 2020, under the TEAMING Grant agreement No. 739572 – The Discoveries CTR. M. G. F. acknowledges the European Research Council under the European Union's Seventh Framework Programme (FP7/2007–2013)/ERC Grant 337753.

References

- 1 S. Datta, L. R. Christena and Y. R. S. Rajaram, *3 Biotech.*, 2013, **3**, 1–9.
- 2 S. C. Silvério, O. Rodríguez, A. P. M. Tavares, J. A. Teixeira and E. A. Macedo, *J. Mol. Catal. B: Enzym.*, 2013, **87**, 37–43.
- 3 A. Kunamneni, S. Camarero, C. García-Burgos, F. J. Plou, A. Ballesteros and M. Alcalde, *Microb. Cell Fact.*, 2008, **7**, 32–47.
- 4 R. A. Sheldon, *Green Chem.*, 2017, **19**, 18–43.
- 5 M. Misson, H. Zhang and B. Jin, *J. R. Soc., Interface*, 2015, **12**, 1–20.
- 6 M. Fernández-Fernández, M. Á. Sanromán and D. Moldes, *Biotechnol. Adv.*, 2013, **31**, 1808–1825.
- 7 Y. Zhang, J. Ge and Z. Liu, *ACS Catal.*, 2015, **5**, 4503–4513.
- 8 A. K. Vahidi, Y. Yang, T. P. N. Ngo and Z. Li, *ACS Catal.*, 2015, **5**, 3157–3161.
- 9 E. Andersson and B. Hahn-Hägerdal, *Enzyme Microb. Technol.*, 1990, **12**, 242–254.
- 10 A. Akelah, in *Functionalized Polymeric Materials in Agriculture and the Food Industry*, ed. A. Akelah, Springer US, Boston, MA, 2013, pp. 195–248, DOI: 10.1007/978-1-4614-7061-8_4.
- 11 M. Martínez-Aragón, S. Burghoff, E. L. V. Goetheer and A. B. de Haan, *Sep. Purif. Technol.*, 2009, **65**, 65–72.
- 12 M. G. Freire, A. F. M. Cláudio, J. M. Araujo, J. A. P. Coutinho, I. M. Marrucho, J. N. Canongia Lopes and L. P. Rebelo, *Chem. Soc. Rev.*, 2012, **41**, 4966–4995.
- 13 M. Martínez-Aragón, S. Burghoff, E. L. V. Goetheer and A. B. de Haan, *Sep. Purif. Technol.*, 2009, **65**, 65–72.
- 14 P.-A. Albertsson, *Nature*, 1958, **182**, 709–711.
- 15 K. E. Gutowski, G. A. Broker, H. D. Willauer, J. G. Huddleston, R. P. Swatloski, J. D. Holbrey and R. D. Rogers, *J. Am. Chem. Soc.*, 2003, **125**, 6632–6633.
- 16 F. A. S. Cunha, A. S. G. Pereira, J. P. A. Fernandes, W. S. Lyra, M. C. U. Araújo and L. F. Almeida, *Food Anal. Methods*, 2017, **10**, 921–930.
- 17 Y. Kohno and H. Ohno, *Chem. Commun.*, 2012, **48**, 7119–7130.
- 18 P. G. Jessop, S. M. Mercer and D. J. Heldebrant, *Energy Environ. Sci.*, 2012, **5**, 7240–7253.

- 19 H. Passos, A. Luís, J. A. P. Coutinho and M. G. Freire, *Sci. Rep.*, 2016, **6**, 20276.
- 20 A. M. Ferreira, A. F. M. Cláudio, M. Valega, F. M. J. Domingues, A. J. D. Silvestre, R. D. Rogers, J. A. P. Coutinho and M. G. Freire, *Green Chem.*, 2017, **19**, 2768–2773.
- 21 A. M. Ferreira, H. Passos, A. Okafuji, M. G. Freire, J. A. P. Coutinho and H. Ohno, *Green Chem.*, 2017, **19**, 4012–4016.
- 22 S. Saita, Y. Mieno, Y. Kohno and H. Ohno, *Chem. Commun.*, 2014, **50**, 15450–15452.
- 23 M. Iqbal, Y. Tao, S. Xie, Y. Zhu, D. Chen, X. Wang, L. Huang, D. Peng, A. Sattar, M. A. B. Shabbir, H. I. Hussain, S. Ahmed and Z. Yuan, *Biol. Proced. Online*, 2016, **18**, 1–18.
- 24 A. Pitto-Barry and N. P. E. Barry, *Polym. Chem.*, 2014, **5**, 3291–3297.
- 25 R. B. Greenwald, *J. Controlled Release*, 2001, **74**, 159–171.
- 26 T. Itoh, *Chem. Rev.*, 2017, **117**, 10567–10607.
- 27 J. F. Pereira, K. A. Kurnia, O. A. Cojocararu, G. Gurau, L. P. Rebelo, R. D. Rogers, M. G. Freire and J. A. P. Coutinho, *Phys. Chem. Chem. Phys.*, 2014, **16**, 5723–5731.
- 28 J. F. Pereira, L. P. Rebelo, R. D. Rogers, J. A. P. Coutinho and M. G. Freire, *Phys. Chem. Chem. Phys.*, 2013, **15**, 19580–19583.
- 29 Y.-M. Lu, Y.-Z. Yang, X.-D. Zhao and C.-B. Xia, *Food Bioprod. Process.*, 2010, **88**, 40–46.
- 30 M. G. Freire, J. F. Pereira, M. Francisco, H. Rodriguez, L. P. Rebelo, R. D. Rogers and J. A. P. Coutinho, *Chem. – Eur. J.*, 2012, **18**, 1831–1839.
- 31 S. Z. Mazlan and S. A. Hanifah, *Int. J. Polym. Sci.*, 2017, **2017**, 1–8.
- 32 L. I. Ramírez-Cavazos, C. Junghanns, N. Ornelas-Soto, D. L. Cárdenas-Chávez, C. Hernández-Luna, P. Demarche, E. Enaud, R. García-Morales, S. N. Agathos and R. Parra, *J. Mol. Catal. B: Enzym.*, 2014, **108**, 32–42.
- 33 H. D. Willauer, J. G. Huddleston, M. Li and R. D. Rogers, *J. Chromatogr. B: Biomed. Sci. Appl.*, 2000, **743**, 127–135.
- 34 H. Walter and E. J. Krob, *J. Chromatogr., A*, 1988, **441**, 261–273.
- 35 A. F. M. Cláudio, J. F. B. Pereira, P. D. McCrary, M. G. Freire, J. A. P. Coutinho and R. D. Rogers, *Phys. Chem. Chem. Phys.*, 2016, **18**, 30009–30019.
- 36 G. Gramss, *Fermentation*, 2017, **3**, 27.
- 37 Chemspider, The free chemical database, <http://www.chemspider.com>, (accessed October 2017).
- 38 E. Nazaruk, A. Michota, J. Bukowska, S. Shleev, L. Gorton and R. Bilewicz, *J. Biol. Inorg. Chem.*, 2007, **12**, 335–344.
- 39 S. Sitthisak, K. Howieson, C. Amezola and R. K. Jayaswal, *Appl. Environ. Microbiol.*, 2005, **71**, 5650–5653.
- 40 M. V. Quental, M. Caban, M. M. Pereira, P. Stepnowski, J. A. P. Coutinho and M. G. Freire, *Biotechnol. J.*, 2015, **10**, 1457–1466.
- 41 D. Mondal, M. Sharma, M. V. Quental, A. P. M. Tavares, K. Prasad and M. G. Freire, *Green Chem.*, 2016, **18**, 6071–6081.
- 42 G. Lugaro, G. Carrea, P. Cremonesi, M. M. Casellato and E. Antonini, *Arch. Biochem. Biophys.*, 1973, **159**, 1–6.
- 43 S. Nicotra, A. Intra, G. Ottolina, S. Riva and B. Danieli, *Tetrahedron: Asymmetry*, 2004, **15**, 2927–2931.
- 44 D. Monti, G. Ottolina, G. Carrea and S. Riva, *Chem. Rev.*, 2011, **111**, 4111–4140.