




Universidade de Aveiro

Departamento de Química



Extraction of Phenolic Compounds with Aqueous Two-Phase Systems

Trabalho realizado por: Ana Filipa Cláudio

Orientação: Prof. João Coutinho e Dra. Mara Freire

Aveiro, 2010

1. Introduction

- 1.1. Scopes and Objectives
- 1.2. Aqueous Two-Phase Systems (ATPS)
- 1.3. Ionic Liquids (ILs)
- 1.4. Phenolic Compounds (PhCs)

2. Extraction of vanillin in ATPS with ILs

- 2.1. Vanillin
- 2.2. Experimental Section
- 2.3. Results and Discussion
- 2.4. Conclusions

3. Extraction of gallic acid in ATPS with ILs

- 3.1. Gallic acid
- 3.2. Experimental Section
- 3.3. Results and Discussion
- 3.4. Conclusions

4. Future Work

1. Introduction

1.1. Scopes and Objectives:

- ❖ Use of aqueous two phase systems (ATPS) composed by ionic liquids (ILs) and typical inorganic salts to develop more benign extraction techniques than those used nowadays;
- ❖ ILs appear as potential and alternative replacements for VOCs due to their negligible vapor pressures and intrinsic character of “designer solvents”;
- ❖ PhCs present general attractive properties, such as their antioxidant, anti-inflammatory, anti-microbial and anticarcinogenic capacities, among others.

1. Introduction

1.2. Aqueous Two-Phase Systems (ATPS)

- ❖ Consist in two aqueous-rich phases mutually incompatible, containing polymer/polymer, polymer/salt or salt/salt combinations;
- ❖ Simple method and easiness in scale-up operations;
- ❖ Used to recover and separate a wide range of biomolecules;
- ❖ The partitioning of the biomolecule depends on the biomolecule size, surface properties, molecular weight, temperature, pH, type of inorganic salts, type of interactions involved, among others.




1. Introduction

1.3. Ionic Liquids (ILs)

Characteristics

Constituted by large and organic cations and organic or inorganic anions;
Melting point below 100 °C;
Negligible flammability and vapour pressure;
High chemical stability in a wide range of temperature;
Easiness in recovery and recycling them;
Most of ILs don't denature biomolecules;
"Benign solvents";
Tunable solvents → "Designer solvents"







Applications

Catalysis;
Organic synthesis;
Chemical reactions;
Chromatographic separations;
Extraction of metal ions;
Separation of biomolecules.

1. Introduction

1.4. Phenolic Compounds (PhCs)

- Present in wood, fruits, vegetables and residues from the industrial or agricultural activities;
- PhCs have a special properties, such as:
 - High antioxidant capacity;
 - Ability of lowering cholesterol;
 - Depression of hypertension;
 - Protection against cardiovascular disease and human leukemia cells;
 - Toxic to bacteria.




2. Extraction of vanillin in ATPS with ILs

2.1. Vanillin

Major compound of vanilla

Used in foods, beverages, pharmaceutical products and fragrance industry.



$pK_a = 8.2$ at 298 K





Due to the scarcity of vanillin in natural sources, the synthesis and recovery of this component is very important.



2. Extraction of vanillin in ATPS with ILs

2.2. Experimental Section


IL
+
 K_3PO_4
+
vanillin



$$K_{\text{Van}} = \frac{[\text{Van}]_{\text{IL}}}{[\text{Van}]_{K_3PO_4}}$$


2.3. Results and discussion

Effect of IL Cations in Vanillin Partitioning at 298.15 K



↑ Alkyl side chains \rightarrow ↓ Surface tension \rightarrow ↑ K_{Van}

↑↑ Alkyl side chains \rightarrow ↑ Dispersive interactions \rightarrow ↓ K_{Van}





Additional double bonds, aromatic or hydroxyl groups

↓
↑ Affinity for water \rightarrow Do not improve the vanillin extraction

❖ The cation-anion interaction strengths are the major forces driving the partitioning of vanillin.

2.3. Results and discussion

Effect of IL Anions in Vanillin Partitioning at 298.15 K



❖ Vanillin partitions preferentially for IL-rich phases composed by Cl^- , Br^- , or $[N(CN)_2]^-$.

❖ The fluorination of $[C_4mim][CH_3SO_3]$ to give $[C_4mim][CF_3SO_3]$


↓
↑ Hydrophobicity → ↑ K_{Van}

❖ $[CH_3SO_3]^-$ and $[CH_3CO_2]^-$ are strongly salting-out inducing ions

↓
High charge density ions
↓
↓ K_{Van}

2.3. Results and discussion

Effect of Temperature in Vanillin Partitioning



❖ Temperature greatly influences the vanillin partitioning



Maximum values in K_{Van}



Energetic and entropic contributions

2.3. Results and discussion

Thermodynamic parameters of transfer - Van't Hoff approach



❖ $\ln(K_{van})$ versus T^{-1} exhibits linearity

$\Delta_{tr}G_m^0 < 0 \rightarrow$ Spontaneous process

$$\ln(K_{van}) = -\frac{\Delta_{tr}H_m^0}{R} \times \frac{1}{T} + \frac{\Delta_{tr}S_m^0}{R}$$

$$\Delta_{tr}G_m^0 = \Delta_{tr}H_m^0 - T\Delta_{tr}S_m^0$$

$\Delta_{tr}H_m^0 < 0 \rightarrow$ Exothermic process

$$\Delta_{tr}G_m^0 = -RT \ln(K_{van})$$




Favourable vanillin-IL interactions

System	$\frac{\Delta_{tr}H_m^0}{kJ \cdot mol^{-1}}$	$\frac{\Delta_{tr}S_m^0}{J \cdot mol^{-1} \cdot K^{-1}}$	$\frac{\Delta_{tr}G_m^0}{kJ \cdot mol^{-1}}$	$\ln(K_{van})$
[C ₄ mim]Cl + K ₃ PO ₄ + water	-23.66	-46.51	-9.79	3.95
[C ₄ mim][CH ₃ SO ₄] + K ₃ PO ₄ + water	-12.62	-15.93	-7.87	3.18
[C ₇ H ₇ mim]Cl + K ₃ PO ₄ + water	-13.19	-13.47	-9.17	3.70
[amim]Cl + K ₃ PO ₄ + water	-24.09	-51.30	-8.80	3.55

$\Delta_{tr}H_m^0$ Depends on anion nature and more complex cations

2.3. Results and discussion

Effect of Concentration in Vanillin Partitioning




↑ [Van] → ↑ K_{Van}

❖ IL cation influences the dependency of the partitioning of vanillin with the solute content.

2.3. Results and discussion

Viscosity of Systems

IL-rich phase (full symbols) and K_3PO_4 -rich phase (open symbols)



3. Extraction of gallic acid in ATPS with ILs

3.1. Gallic acid


Characterists: antioxidant, anti-inflammatory, antifungal, anti-tumor, diuretic, depurative, intestinal antiseptic, bacteriostatic, bactericidal and anti-arthritic.

Pharmaceutical, nutraceutical and cosmetic applications



Treatment of gastric tonus problems, anorexia, bloating, gases, urinary diseases gout, skin repairer and as sedative.

Gallic acid is present in fruits (grapes), pomegranate husk, vegetables, green and black teas, oak wood and residual waste.





$pK_a = 4.4$ at 298 K



3. Extraction of gallic acid in ATPS with ILs

3.2 Experimental Section: Phase Diagrams



▲ [C₈py][N(CN)₂] ♦ [C₄mim][CF₃SO₃] ◆ [C₄mim][TOS] ▲ [C₄mim][N(CN)₂] ● [C₇H₇mim][C₂H₅SO₄] × [C₂mim][CF₃SO₃]
◆ [C₄mim][C₂H₅SO₄] ◆ [C₄mim][CH₃SO₄] ■ [C₄mim]Br + [C₇H₇mim]Cl - [C₈mim]Cl ■ [C₇mim]Cl



Anions or cations with more hydrophobic characteristics have a greater capacity to promote ATPS



ILs containing the anions [N(CN)₂]⁻ and [CF₃SO₃]⁻ or larger alkyl side chains at the cation

3. Extraction of gallic acid in ATPS with ILs

Phase Diagrams



❖ Fixing [C₄mim]⁺



[CF₃SO₃]⁻ > [TOS]⁻ > [N(CN)₂]⁻
> [C₂H₅SO₄]⁻ > [CH₃SO₄]⁻ > Br⁻

❖ Fixing [C₇H₇mim]⁺



[C₂H₅SO₄]⁻ > Cl⁻

❖ ↑ alkyl chain in anion or cation



↑ ability of the IL for ATPS formation

❖ Higher capability by ILs to promote ATPS



with stronger salting-in inducing behavior

❖ longer alkyl chains




↑ ability of the IL for ATPS formation

3.3. Results and discussion



Effect of IL ions in the acid gallic partitioning

↑ Alkyl chains cation or anion → ↓ Surface tension → ↑ K_{GA}
↑↑ Alkyl side chains → ↑ Dispersive interactions → ↓ K_{GA}



3.3. Results and discussion

Effect of pH in the acid gallic partitioning



IL+inorganic salt + water		Na ₂ SO ₄	K ₂ HPO ₄ /KH ₂ PO ₄	K ₃ PO ₄
[C ₂ mim][CF ₃ SO ₃]	Salt-rich phase	3.32	7.28	13.09
	IL-rich phase	2.71	7.57	13.15
[C ₄ mim][CF ₃ SO ₃]	Salt-rich phase	3.04	7.10	12.85
	IL-rich phase	3.12	7.37	13.10
[C ₇ mim]Cl	Salt-rich phase	4.16	7.22	12.85
	IL-rich phase	4.15	7.45	12.99

❖ pH at the salt-rich phase ≈ pH at the IL-rich phase for the same system;

❖ The cation alkyl chain length does not significantly contribute for differences in pH;

❖ Changing the anion from [CF₃SO₃]⁻ to Cl⁻ guides to slightly differences in the acidity of both aqueous phases.

4. Future Work

- ❖ Extend this type of study to other phenolic compounds and test more conditions in order to optimize extraction routes;
- ❖ Determine the pH of both rich phases for some systems already evaluated in this work;
- ❖ In order to better evaluate the effect of pH on the phenolic compounds extraction it will be important to extend the study for more systems based on different ILs;
- ❖ Use the knowledge acquired to proceed to more practical and real experiments attempting direct extractions from biomass, such as the extraction of phenolic compounds from wood, plants or wastewater effluents.

Thank you for your attention!