



Improvements in the enzymatic degradation of textile dyes using ionic-liquid-based surfactants

Rui M.F. Bento, Mafalda R. Almeida, Pankaj Bharmoria, Mara G. Freire, Ana P.M. Tavares*

CICECO – Aveiro Institute of Materials, Department of Chemistry, University of Aveiro, 3810-193 Aveiro, Portugal

ARTICLE INFO

Keywords:

Laccase
Enzymatic activity
Surfactant-based ionic liquids
Indigo carmine
Textile aqueous effluents

ABSTRACT

The intensive use of water containing dyes by the textile industry, and consequently the contamination of soils and water, represents serious environmental concerns. Amongst the several processes applied in the treatment of textile effluents, biological-based processes, if designed to be cost-effective and ecofriendly, are promising alternatives to decolorize textile effluents. In this work we investigate and propose the novel use of ionic liquids (ILs) with surfactant characteristics to improve the degradation of the largely used and highly hydrophobic textile dye indigo carmine (IC) by laccase. An initial screening on the activity of laccase in aqueous solutions of twelve surfactant-based ILs from three different families, namely tetraalkylammonium- and imidazolium-based cationic surfactants and cholinium-based anionic surfactants, at different concentrations, was carried out. Significant improvements in the activity of laccase were observed with decyltrimethylammonium bromide, $[N_{10111}]Br$, and 1-decyl-3-methylimidazolium chloride, $[C_{10}mim]Cl$, at 75 mM (above the critical micellar concentration of each IL). These ILs were then investigated in aqueous solutions to simultaneously encapsulate laccase and IC for the *in situ* enzymatic biodegradation of the dye. The use of ILs remarkably increases the degradation rate of the dye and decolorization efficiency; a degradation efficiency of IC of 82% is attained in 0.5 h using aqueous solutions of $[N_{10111}]Br$, whereas without IL only 6% of IC is degraded. Furthermore, 93% of the dye decolorization was achieved with $[N_{10111}]Br$. The overall gathered results show that it is possible to significantly improve the degradation of hydrophobic dyes by enzymes using appropriate surfactant-based ILs, while foreseeing the use of the treated water by the same textile industries in new dyeing steps and thus contributing to a substantial decrease of the economic input and environmental footprint of these industries.

1. Introduction

Serious environmental concerns have arisen due to industrial developments, leading to significant risks to human health and to the ecosystem [1]. In particular, the textile industry plays a pivotal role contributing to these concerns. Textiles production requires several stages of mechanical processing, involving the discharge of a wide variety of pollutants such as textile dyes [2]. Furthermore, the textile industry is highly water intensive, requiring about 200 L of water to produce 1 kg of textiles [3,4]. The textile effluents are enriched in dyes used in the dyeing process, many of them toxic and mutagenic [5]. Synthetic dyes, such as indigo carmine (IC) – a typical recalcitrant dye, are highly persistent and their removal from wastewater is difficult since they are designed to be chemically and photolytically stable [6]. Accordingly, the removal and degradation of dyes is a challenging issue from the environmental perspective because conventional methods for the treatment of aqueous systems, such as ozonation, coagulation-flocculation, oxidation, precipitation, adsorption and ion

exchange are not totally effective in dyes removal [1,7,8]. Therefore, the development of new strategies for the removal or degradation of dyes from textile effluents, ideally allowing the water recycling by the same textile industry, is in high demand.

Biological-based treatment technologies, if designed to be cost-effective and ecofriendly technologies, are promising alternatives to decolorize textile effluents [9]. Bioremediation using enzymes have gained significant notoriety due to its versatility and efficiency in the degradation of persistent organic pollutants from wastewater [10,11]. Oxidative enzymes such as laccases, peroxidases and tyrosinases have high potential in the oxidation of persistent environmental pollutants [12]. These enzymes have the capacity to convert the target pollutants into less toxic or insoluble compounds, which can be then removed from effluents [13]. Laccase (benzenodiol oxygen oxidoreductase, EC 1.10.3.2) is an efficient multicopper oxidase exhibiting a broad substrate specificity, e.g. phenols and aromatic amines, being thus used in industrial, biotechnological and environmental applications [14,15]. More specifically, its ability to

* Corresponding author.

E-mail address: aptavares@ua.pt (A.P.M. Tavares).

<https://doi.org/10.1016/j.seppur.2019.116191>

Received 3 June 2019; Received in revised form 29 September 2019; Accepted 7 October 2019

Available online 08 October 2019

1383-5866/ © 2019 Elsevier B.V. All rights reserved.

degrade a variety of dyes [16,17], such as IC largely used by the textile industry, leads to its use in bioremediation processes with the aim of reducing the environmental impact caused by textile industrial effluents [18]. However, the efficient application of enzymes in industrial processes is not always successful due to the enzymes labile nature and loss of stability and enzymatic activity [19].

The use of micellar systems and emulsions can improve catalytic reactions due to the enzyme superactivity phenomenon, being micellar enzymology a hot topic of research [20]. In addition to conventional and largely used surfactants, ionic liquids (ILs) with surfactant properties can be designed and applied in the micellar enzymology field. ILs have gained particular importance in the field of biocatalysis and are currently recognized as promising solvent media [21,22]. ILs generally exhibit large organic cations and a variety of anions that can be organic or inorganic, which leads to a decrease of these salts melting temperatures when compared to conventional salts [23,24]. Being organic salts, most aprotic ILs display negligible vapor pressures, low flammability, and high thermal and chemical stability [25]. Furthermore, due to their tunable character achieved by altering the anion or cation chemical structure, it is possible to tailor the properties of ILs and to synthesize a specific IL to a target reaction or application. Accordingly, long alkyl side chain ILs, if properly designed, may be amphiphilic compounds able to self-aggregate and form micelles [26,27]. A wide range of ILs has been characterized in terms of their critical micellar concentration (CMC), being investigated in several applications as alternative surfactants or as co-surfactants [28,29].

Besides the large number of studies demonstrating the ability of ILs to form micelles and determination of their CMC values, fewer works addressed the use of ILs with surfactant characteristics to improve enzymatic bioreactions [28,29]. In this work, we investigate and propose the use of ILs with surfactant behavior to improve the degradation of the IC dye by laccase, which may allow the reuse of the discharged water by the textile industry.

2. Experimental

2.1. Chemicals

Laccase from *Trametes versicolor* (≥ 10 U mg⁻¹) was purchased from Sigma-Aldrich. Three families of ILs were investigated, namely 1-alkyl-3-methylimidazolium chloride and 1-alkyl-trimethylammonium bromide (acting as cationic surfactants), and cholinium carboxylates (acting as anionic surfactants). The cationic surfactant-based ILs 1-methyl-3-octylimidazolium chloride (> 98% purity), 1-decyl-3-methylimidazolium chloride (> 98% purity), 1-dodecyl-3-methylimidazolium chloride (> 98% purity), 1-methyl-3-tetradecylmethylimidazolium chloride (> 98% purity), were purchased from Iolitec. Octyltrimethylammonium bromide (98% purity) and decyltrimethylammonium bromide (99% purity) were acquired from TCI Europe N.V. Dodecyltrimethylammonium bromide (99% purity) and tetradecyltrimethylammonium bromide (98% purity) were supplied by Alfa Aesar. The cholinium-based carboxylates, namely cholinium octanoate, cholinium decanoate, cholinium dodecanoate and cholinium tetradecanoate were synthesized by us according to the protocol described in the literature [30]. The abbreviation, molecular weight and critical micellar concentration (CMC) of all ILs are given in Table 1, and the ILs chemical structures are shown in Fig. 1. The enzyme substrate 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt (ABTS) was obtained from Sigma-Aldrich. Indigo carmine (IC) (C.I. 73015) was supplied by ACROS Organics™. The chemical structure and properties of IC are provided in Table 2. All aqueous solutions were prepared using Milli-Q ultrapure water.

2.2. Laccase activity in aqueous solutions of surfactant-based ILs

An initial screening on potential ILs to improve the laccase activity was carried out. To this end, aqueous solutions containing laccase and ILs were prepared with a final enzyme activity of 1000 U L⁻¹. IL concentrations of

10, 50, 100, 250 and 350 mM were used, and assays were performed at room temperature (ca. 25 °C) and pH 4.5 (adjusted with HCl 0.5 M). A control solution was used, with no addition of IL, at the same experimental conditions. The laccase activity in presence of ILs was determined spectrophotometrically by monitoring the oxidation of ABTS at 420 nm ($\epsilon = 36,000$ M⁻¹cm⁻¹), using a SHIMADZU UV-1800, UV-Vis Spectrophotometer. The enzymatic reaction was carried out by adding 50 μ L of sample in 250 μ L of ABTS 1.6 M and 700 μ L of Milli-Q water (adjusted at pH 4.5 with HCl 0.5 M). One unit (U) of laccase activity is defined as the amount of enzyme that oxidized 1 μ mol of ABTS (molar extinction coefficient [ϵ_{420}], 36,000 M⁻¹cm⁻¹) per minute. The laccase activity is presented in U L⁻¹. All assays were repeated at least three times.

2.3. Enzymatic degradation of IC in ILs aqueous solutions

Surfactant-based ILs in which the enzyme showed higher activities were selected to study the enzymatic degradation of IC. The decolorization of IC was evaluated by the monitorization of the absorbance at 608 nm using a SHIMADZU UV-1800, UV-VIS Spectrophotometer. The enzymatic reaction mixture contained: 2.50 mL of Milli-Q water at pH 4.5, 1.25 mL of IC (25 mg L⁻¹), 1.25 mL of laccase (1000 U L⁻¹) and IL (20 and 75 mM, below and above the CMC of the ILs used in this step, respectively). Control aqueous solutions composed of each IL and of IC were used. At the same conditions, no dye degradation was observed with these solutions, showing that laccase is responsible for the dye degradation.

The decolorization of IC extent was determined according to equation (1):

$$\text{IC decolorization (\%)} = \left(\frac{A_0}{A_t} \right) \times 100 \quad (1)$$

where A_0 and A_t are the initial absorbance and the absorbance observed at a certain time (t), respectively. All assays were performed in triplicate.

2.4. Circular dichroism spectroscopy

The laccase secondary structure in presence of [N₁₀₁₁₁]Br and [C_{10mim}]Cl was evaluated by circular dichroism (CD) spectroscopy using a Jasco J-1500CD spectrometer. Aqueous solutions containing laccase and [N₁₀₁₁₁]Br or [C_{10mim}]Cl were prepared with a final enzyme concentration of 0.6 mg mL⁻¹. We could not use high concentrations of ILs on ground of the fact that the studied ILs show high absorbance in the far UV region beyond 1 mM. "Blank" solutions at the same IL concentrations (with no laccase added) were used to remove the ILs interference on the CD spectrum. A control using aqueous laccase solution was used. CD spectra were recorded from 190 to 260 nm using quartz CD cuvettes (0.1 cm) at room temperature. Each CD spectrum is the result of five accumulations recorded in millidegrees. The following acquisition parameters were used: data pitch, 1.0 nm; sensitivity 100 mdeg; response time 4 s; bandwidth, 0.50 nm; and scan speed, 100 nm min⁻¹. A smooth was performed in each CD spectrum using the following parameters: method, Savitzky-Golay; convolution width, 7.

2.5. Analysis by optical microscopy

To evaluate the microscopic appearance of the micellar [N₁₀₁₁₁]Br solutions, different samples were prepared: IL (75 mM) and IC (25 mg L⁻¹); IL (75 mM) and laccase (1000 U L⁻¹) and IL (75 mM), laccase (1000 U L⁻¹) and IC (25 mg L⁻¹). The solutions were centrifuged (microfuge Star 17, VWR) at increasing speeds from 300 to 13300 rpm attempting the aggregates precipitation. Microscopic images of the precipitated fraction were obtained using a polarized light microscope, Olympus BX51 with 100 \times and 200 \times magnifications.

Table 1

Name, abbreviation, molecular weight and critical micellar concentration (CMC) of the surfactant-based ionic liquids (ILs) studied.

IL	Abbreviation	Molecular weight (g mol^{-1})	CMC (mM) [31–33]
Octyltrimethylammonium bromide	[N ₈₁₁₁]Br	252.23	130.0
Decyltrimethylammonium bromide	[N ₁₀₁₁₁]Br	280.29	64.6
Dodecyltrimethylammonium bromide	[N ₁₂₁₁₁]Br	308.34	15.6
Tetradecyltrimethylammonium bromide	[N ₁₄₁₁₁]Br	336.39	3.6
1-Methyl-3-octylimidazolium chloride	[C ₈ mim]Cl	230.78	233.0
1-Decyl-3-methylimidazolium chloride	[C ₁₀ mim]Cl	258.83	58.7
1-Dodecyl-3-methylimidazolium chloride	[C ₁₂ mim]Cl	286.88	15.2
1-Methyl-3-tetradecylmethylimidazolium chloride	[C ₁₄ mim]Cl	314.94	3.9
Cholinium octanoate	[Ch][C ₈ O ₂]	247.37	303.3
Cholinium decanoate	[Ch][C ₁₀ O ₂]	275.43	104.3
Cholinium dodecanoate	[Ch][C ₁₂ O ₂]	303.48	25.8
Cholinium tetradecanoate	[Ch][C ₁₄ O ₂]	331.53	7.0

3. Results and discussion

3.1. Laccase activity in aqueous solutions of surfactant-based ILs

The development of the micellar enzymology concept, applying surfactant-based ILs, has been a hot topic of research due to the enzymes superactivity phenomenon afforded by amphiphilic ILs [20]. However, the role of the IL chemical structure and properties on the laccase activity and further applications is still scarce. Thus, it is of high importance to identify appropriate IL-based surfactants able to improve the catalytic activity of laccases and related bioreactions performance.

Since enzyme-catalyzed reactions are strongly influenced by the reaction solvent medium, the chemical structure and concentration of a series of IL-based surfactants were first evaluated toward the laccase activity. Aqueous solutions with laccase at a final concentration of enzyme of $\sim 1000 \text{ U L}^{-1}$ and ILs with concentrations of 10, 50, 100,

250 and 350 mM were investigated. These concentrations were selected to be below and above the CMC of each IL (Table 1). Twelve surfactant-based ILs from three different families, namely tetraalkylammonium- and imidazolium-based cationic surfactants and cholinium-based anionic surfactants were investigated. A control solution (with no IL at the same experimental conditions – pH 4.5 adjusted with HCl 0.5 M) was prepared and always used for comparison purposes. The relative laccase activity in the different IL aqueous solutions is depicted in Fig. 2, whose detailed results are provided in Table S1 in the Supporting Information. The relative laccase activity corresponds to the percentage activity of laccase in the control in respect to each IL aqueous solution.

The results obtained show that reactions catalyzed by laccase are considerably influenced by the IL chemical structure, i.e. if it is an anionic or cationic surfactant, and by the IL concentration. In general, the laccase activity in presence of the cationic tetraalkylammonium- and imidazolium-based surfactants is higher at lower IL concentrations





Fig.1. Chemical structures of the surfactant-based ionic liquids investigated.

Table 2
Structure and properties of the indigo carmine (IC) dye used in the micellar enzymatic degradation tests [34].

Dye	Molecular weight (g mol^{-1})	Water solubility at 25 °C (g L^{-1})	Log K_{ow}	Chemical structure
Indigo carmine (IC)	466.3	10.0	-0.991	




Fig. 2. Relative laccase activity (%) in aqueous solutions composed of (A) ammonium-based ionic liquids (ILs); (B) imidazolium-based ILs; and (C) cholinium-based ILs. Arrows indicate the critical micellar concentration (CMC) of each surfactant-based IL.

and with ILs with shorter alkyl side chains, reaching values similar to the control (> 85% of relative activity) with the following ILs at the following concentrations: [N₈₁₁₁]Br at 10 and 50 mM; [N₁₀₁₁₁]Br at 10, 50 and 100 mM; [C₈mim]Cl at 10, 50 and 100 mM; [C₁₀mim]Cl at 10, 50 and 100 mM; [C₁₂mim]Cl at 10 mM; and [C₁₄mim]Cl at 10 mM. The dependence of the laccase activity with the alkyl chain length can be attributed to hydrophobic interactions occurring between the IL and the hydrophobic moieties of the enzyme which can affect the enzyme conformational structure and consequently its reactivity [35,36]. Among these, [C₁₀mim]Cl and [N₁₀₁₁₁]Br at 75 mM are above the respective CMC values (Table 1), meaning that micelles are present. On

the other hand, the laccase activity in presence of cholinium-based ILs (anionic surfactants) is not favored; in fact, a maximum relative activity of 50% has been observed with these ILs when compared to the control. Furthermore, the laccase activity decreases with the increase in the IL concentration. These results indicate that the studied anionic surfactants act as inhibitors of laccase. IL ions can interact with the enzyme by electrostatic interactions, occurring between the surfactant head groups and charged amino acid residues, and by hydrophobic forces occurring between the IL alkyl chains and the enzyme hydrophobic cores [37]. Moreover, other physicochemical properties of ILs such as polarity, water miscibility, viscosity, and ions charge density may also influence enzymatic activity, and have been studied [38].

In the literature, it is reported that for anionic surfactants, the initial binding between the enzyme and the surfactant occurs on the cationic sites of the enzyme (amino acids side chains) causing the unfolding (denaturation) of the enzyme and the formation of a strong protein-surfactant complex [37]. These reported evidences suggest that a similar phenomenon may happen in the current work when addressing anionic IL surfactants. A recent review from Liu et al. [39] describes the ILs features that influence the laccase activity. For instance, the authors stated that there is no correlation between the IL polarity and the laccase activity [39]. On the other hand, Yang et al. [40] described that the effect of the ILs' ions on the enzyme stability be explained by the Hofmeister series. Overall, the authors concluded that chaotropic cations and kosmotropic anions stabilize the enzyme, whereas kosmotropic cations and chaotropic anions tend to destabilize it [39].

For our results, it is clear that the catalytic activity of laccase is affected by the IL type and concentration. Taking into account the overall results and the foreseen application of using micellar systems combined with laccase to degrade highly hydrophobic textile dyes, the ILs [N₁₀₁₁₁]Br (CMC = 64.6 mM) and [C₁₀mim]Cl (CMC = 58.7 mM) with a high activity of laccase (> 85%) at an IL concentration of 20 and 75 mM were selected and investigated in the following set of studies regarding the IC decolorization.

3.2. Enzymatic degradation of IC in ILs aqueous solutions

The mechanism of IC degradation by laccase was studied by Campos et al. [41]. Laccase induces the IC oxidation, resulting in the formation of the indole-2,3-dione intermediate, which is further decomposed into 2-aminobenzoic acid [41]. This reaction occurs due to the trinuclear copper centers of laccase that oxidize the dye, transferring the electrons from the substrate to the copper center, while O₂ is reduced to water [42]. The ecotoxicological features of IC reaction products were evaluated by MicroTox, in which it was demonstrated that they are less toxic than the IC precursor [43]. The toxicity of IC degradation products by an enzyme preparation (not defined) was addressed against fish (*Talapia*) and two bacteria (*Escherichia coli* and *Bacillus subtilis*), with no toxicity obtained [44]. Furthermore, toxicological studies of IC using an integrated treatment by a trimeric thermostable laccase and a microbial consortium showed that the degradation products were non-toxic, while the initial IC was toxic [45]. These results support our strategy on resorting to biocatalysis to perform the degradation of dyes aiming water treatment.

Taking into account that micellar systems are required due to the high hydrophobicity (log(K_{ow}): -0.991) and low water solubility of IC (Table 2), the effects afforded by [N₁₀₁₁₁]Br and [C₁₀mim]Cl at 20 and




Fig. 3. (A) Indigo Carmine (IC) decolorization by laccase alone (●), IC decolorization by laccase in [C₁₀mim]Cl (75 mM) aqueous solutions (◆), IC decolorization by laccase in [C₁₀mim]Cl (20 mM) aqueous solutions (◇), IC decolorization by laccase in [N₁₀₁₁₁]Br (75 mM) aqueous solutions (■), and IC decolorization by laccase in [N₁₀₁₁₁]Br (20 mM) aqueous solutions (□) at 25 °C; (B) Images showing the indigo carmine decolorization by laccase in the presence of [N₁₀₁₁₁]Br (75 mM) aqueous solutions at 25 °C.

75 mM (below and above their CMC values) on the color removal of IC by laccase were evaluated. Aqueous solutions of ILs above their CMC will allow the formation of micelles able to encapsulate laccase and IC for the *in situ* enzymatic biodegradation.

Controls composed of each IL and IC (individually) were prepared and no degradation of IC was observed, showing the main role and requirement of laccase in the target oxidative reaction. Although there are several studies on the decolorization of dyes by laccase [46–50], to the best of our knowledge, there are no literature reports on dyes degradation by enzymes promoted by IL-based surfactants. Fig. 3(A) and Table S2 in the Supporting Information provide the results obtained for the decolorization of IC by laccase in the presence and absence of [N₁₀₁₁₁]Br or [C₁₀mim]Cl aqueous solutions. The decolorization of IC by laccase shows a faster and higher catalytic performance in the presence of both surfactant-based ILs at 75 mM when compared to tests performed with 20 mM of IL (below their CMC) and the control (laccase with no IL at the same conditions, Fig. 3(A)), showing the beneficial role of ILs in improving biocatalytic reactions. For instance at 0.5 h of reaction, the values increase from 6% of IC decolorization by laccase with no IL added to 30% and 82% when using the ILs [C₁₀mim]Cl and [N₁₀₁₁₁]Br at 75 mM, respectively. Comparing both surfactant-based ILs, [N₁₀₁₁₁]Br performs better and leads to a higher decolorization efficiency at all times evaluated. Remarkable, after 24 h, an almost complete dye decolorization (> 90%) was achieved, representing an improvement of three times when compared to aqueous solution of ILs at 20 mM and the control with no IL added.

Fig. 3(B) shows the macroscopic appearance regarding the IC decolorization by laccase with 75 mM of [N₁₀₁₁₁]Br. Overall, the obtained results reveal a higher dye decolorization performance by laccase in presence of surfactant-based ILs aqueous solutions. These results do not have the same tendency to the results discussed before, where the laccase activity and the oxidation of ABTS decreases with the increase

in the IL concentration. According to Liu et al. [39], laccase has broad substrate specificities, and the design of general rules for selecting efficient ILs-laccase systems is not a straightforward task. For instance, using the same laccase, the activity towards catechol was partially inhibited by 1-butyl-3-methylimidazolium bromide (at 20% v/v), and completely eliminated when dealing with ABTS [51,52]. Therefore, the enzyme-substrate affinity should be taken into consideration to define the role of IL in laccase-catalysed reactions.

The results of IC decolorization by laccase suggest that by using the respective ILs at concentrations above their CMC, the highly hydrophobic substrate (dye) is inside the micelles as well as the enzyme, turning the dye more accessible to the enzyme in this microenvironment and improving the enzyme catalytic performance. According to the literature, IC cannot be oxidized at high levels by laccase alone. Accordingly, specific laccase mediators, such as 1-hydroxybenzotriazole, 3,5-Dimethoxy-4-hydroxybenzaldehyde, ABTS, violuric acid or N-hydroxyphthalimide [53,54], are usually added to improve the laccase oxidation of IC. In a different approach, IC was degraded by laccase without mediator, but in this case a mutated laccase, with a different active site, was used, allowing to remove a maximum of ~60% of IC [55]. In this work, a fast and efficient degradation of IC without the presence of mediators or enzymes mutation was successfully obtained applying surfactant-based ILs.

To better understand the effect of ILs on the structure-activity relation of laccase the changes in the secondary structure of laccase were appraised by the analysis of the CD spectra. The secondary structure of native laccase shows a characteristic positive peak at around 198 nm (Fig. 4, Fig. S1 in the Supporting Information). Quantitative analysis of the secondary structure was done using K₂D₃ online secondary structural analysis software in the wavelength range from 240 to 190 nm [56,57] and found to be α -helical = 3% and β -sheet = 34%. The calculated secondary structure is in close proximity to that reported for crystal structure of laccase from *Trametes versicolor* (PDB. IGCY; α -helical = 11% and α -helical = 37%), hence demonstrating the accuracy of the spectra acquisition and treatment. Moreover, ionic surfactants have been reported to alter the structure of proteins mainly in the low concentration regime, therefore validating the structural studies at low concentration carried out in this work. At the studied concentration both ILs alter the structure of laccase, although not significantly (Fig. 4, Fig. S1 in the Supporting Information). [N₁₀₁₁₁]Br was found to induce more changes in the structure of laccase as compared to [C₁₀mim]Cl. Comparing the CD spectra and the IC degradation results in the presence of ILs at a concentration below the CMC it can be concluded that the altered structure of laccase is more active due to the exposure of the active sites to a higher number of substrate molecules. Bharmoria et al. [58] have reported a similar behavior of enzyme cellulase upon interaction with the IL 3-methyl-1-octylimidazolium dodecylsulfate at low concentration. The higher activity of laccase in the micellar solution of both ILs can be accounted by the availability of hydrophobic micellar




Fig. 4. CD spectra of aqueous solution of laccase (control), blue line; aqueous solution of laccase and [C₁₀mim]Cl (1 mM), orange line; and aqueous solution of laccase and [N₁₀₁₁₁]Br (1 mM), grey line.

interfaces allowing better interactions between the hydrophobic dye as the substrate and the enzyme in its altered form. In different reports, a similar behavior was observed for the peroxidase activity of cytochrome *c*, which enhanced in the vesicular solution of surfactant IL; cholinium dioctylsulfosuccinate (a surface active ionic liquid) induced a conformational transition in the secondary structure of cytochrome *c* with an enhanced peroxidase activity [59,60].

To evaluate if the current laccase reaction occurs inside or outside the ILs micelles, different mixtures containing $[N_{10111}]Br$, substrate (IC) and laccase were prepared: (i) $[N_{10111}]Br$ and laccase; (ii) $[N_{10111}]Br$ and IC; and (iii) $[N_{10111}]Br$, laccase and IC. After 2.5 h, all solutions were centrifuged in a progressive rotational speed aiming at precipitating the ILs aggregates containing laccase and the substrate. The obtained precipitates were analyzed by optical microscopy. Fig. 5 shows the macroscopic and microscopic appearance of the different $[N_{10111}]Br$ aqueous solutions and respective precipitates. A blue precipitate with the mixture composed of $[N_{10111}]Br$ + IC and a colorless precipitate with the $[N_{10111}]Br$ + IC + enzyme were obtained. From the analysis of the microscopic images shown in (Fig. 5) it seems that IC is mostly confined inside the aggregates formed by the $[N_{10111}]Br$ IL. These macroscopic results suggest that the oxidation of IC takes place inside the micelles, where both the enzyme and the dye are incorporated thus promoting an efficient and improved IC decolorization. Moreover, for this IL, the enzyme activity was measured in the supernatant after a centrifugation step. No enzymatic activity was detected, being one additional indication that the enzyme is inside the aggregates formed by the $[N_{10111}]Br$ IL. These findings are in accordance with Liang et al. [61], who suggested that the inner of micelles is a suitable mimetic environment of living cells, supporting the enzymes “superactivity” in micelles cores.

The overall gathered results show that it is possible to improve the degradation of hydrophobic dyes by enzymes using appropriate surfactant-based ILs. Furthermore, large aggregates are formed, which can be removed by precipitation or filtration, allowing the dyes removal and water treatment. Even if some IL is still present in the treated water,

these have shown to be advantageous in the dyeing of wool, polyester, and cotton with the Disperse Red 13 dye in the absence of auxiliary agents [62]. Accordingly, in the current work, we foresee the use of the treated water by the same textile industries in new dyeing steps, contributing to a significant decrease of the economic input and environmental footprint of these industries.

4. Conclusions

In this work, we investigated and proposed the use of ILs with surfactant behavior to improve the degradation of the IC dye by laccase, which may allow the reuse of the discharged water by the textile industry. The activity of laccase in aqueous solutions of three families of ILs, namely 1-alkyl-3-methylimidazolium chloride ($[C_n\text{mim}][Cl]$) and 1-alkyltrimethylammonium bromide ($[N_n][Br]$) as cationic surfactants and cholinium carboxylate ($[Ch][C_nO_2]$) as anionic surfactants, was evaluated. A high activity of laccase was obtained with $[N_{10111}]Br$ and $[C_{10}\text{mim}][Cl]$ at 75 mM, above their CMC, and where ca. 90% of the enzyme residual activity is maintained. These ILs were then investigated to improve the color removal of IC by laccase. It was demonstrated that the use of the $[N_{10111}]Br$ IL is favorable for the enzymatic degradation of the dye. Remarkably, a significantly higher and fast decolorization of the IC dye was obtained, and within 0.5 h it was possible to achieve a color removal percentage of 82% (against 6% achieved without IL). After 24 h, 93% of the dye decolorization was accomplished in the presence of the same IL at the same concentration. Overall, this work shows the possibility of using surfactant-based ILs instead of the commonly used mediators or enzymes mutation approaches to significantly improve the enzymatic degradation of textile dyes.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

This work was developed within the scope of the project CICECO-Aveiro Institute of Materials, FCT Ref. UID/CTM/50011/2019, financed by national funds through the FCT/MCTES. This work was also financially supported by the project POCI-01-0145-FEER-031106 (IonCytDevice) and POCI-01-0145-FEDER-031268 funded by FEDER, through COMPETE2020 – Programa Operacional Competitividade e Internacionalização (POCI), and by national funds (OE), through FCT/MCTES. Ana P.M. Tavares acknowledges FCT for the Investigator Programme (IF/01634/2015).

Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.seppur.2019.116191>.

References

- [1] A. Shah, S. Shahzad, A. Munir, M.N. Nadagouda, G.S. Khan, D.F. Shams, et al., Micelles as soil and water decontamination agents, *Chem. Rev.* 116 (10) (2016) 6042–6074.
- [2] H. Hayat, Q. Mahmood, A. Pervez, Z.A. Bhatti, S.A. Baig, Comparative decolorization of dyes in textile wastewater using biological and chemical treatment, *Sep. Purif. Technol.* 154 (2015) 149–153.
- [3] S. Vineta, Z. Silvana, R. Sanja, S. Golomeova, Methods for waste waters treatment in textile industry, in: International Scientific Conference – “UNITECH 2014” – Gabrovo, 2014.
- [4] D. De Jager, M.S. Sheldon, W. Edwards, Colour removal from textile wastewater using a pilot-scale dual-stage MBR and subsequent RO system, *Sep. Purif. Technol.* 135 (2014) 135–144.
- [5] F. Orts, A.I. del Río, J. Molina, J. Bonastre FC, Study of the reuse of industrial wastewater after electrochemical treatment of textile effluents without external addition of chloride, *Int. J. Electrochem. Sci.* 14 (2019) 1733–1750.
- [6] L.L. Jiayi Wang, F. Feng, Improving the indigo carmine decolorization ability of a




Fig. 5. Indigo carmine and $[N_{10111}]Br$ aqueous solutions and optical microscope images of the precipitate obtained after centrifugation.

- bacillus amyloliquefaciens laccase by site-directed mutagenesis, *Catalysts* 7 (9) (2017) 275.
- [7] B. Sharma, A.K. Dangi, P. Shukla, Contemporary enzyme based technologies for bioremediation: a review, *J. Environ. Manage.* 210 (2018) 10–22.
- [8] M. Riera-Torres, C. Gutiérrez-Bouzán, Optimisation of the electrochemical and UV combined treatment to remove colour and organic halogenated compounds of textile effluents, *Sep. Purif. Technol.* 98 (2012) 375–382.
- [9] K.D. Mojsov, Biotechnological applications of laccases in the textile industry, *Adv. Technol.* 3 (1) (2014) 76–79.
- [10] N. Durán, E. Esposito, Potential applications of oxidative enzymes and phenoloxidase-like compounds in wastewater and soil treatment: a review, *Appl. Catal. B Environ.* 28 (2) (2000) 83–99.
- [11] C.S.K. Rao, S.S. Rao, Role of microbial enzymes in the bioremediation of pollutants. A review, *Enzyme Res.* 2011 (2011) 1–11.
- [12] A.P.M. Tavares, A.M.B.R. Xavier, Application of oxidative enzymes in the treatment of industrial effluents, *Recent Developments in Biotechnology: Biodegradation and Bioremediation*, Edition: Studium Press LLC, Houston, TX, USA, 2014, pp. 193–214.
- [13] M. Husain, Q. Husain, Applications of redox mediators in the treatment of organic pollutants by using oxidoreductive enzymes: a review, *Crit. Rev. Environ. Sci. Technol.* 38 (1) (2007 Dec) 1–42.
- [14] M. Tisma, P. Znidarsic-Plazl, D. Vasic-Racki, B. Zelic, Optimization of laccase production by *Trametes versicolor* cultivated on industrial waste, *Appl. Biochem. Biotechnol.* 166 (1) (2012) 36–46.
- [15] Adinarayana Kunamneni, F.J.P. Antonio Ballesteros, M. Alcalde, Fungal laccase – a versatile enzyme for biotechnological applications, *Commun. Curr. Res. Educ. Top Trends Appl. Microbiol.* 1 (2007) 233–245.
- [16] H.-M. Xu, X.-F. Sun, S.-Y. Wang, C. Song, S.-G. Wang, Development of laccase/graphene oxide membrane for enhanced synthetic dyes separation and degradation, *Sep. Purif. Technol.* 204 (2018) 255–260.
- [17] S. Moreira, A.M.F. Milagres, S.I. Mussatto, Reactive dyes and textile effluent decolorization by a mediator system of salt-tolerant laccase from *Peniophora cinerea*, *Sep. Purif. Technol.* 135 (2014) 183–189.
- [18] C. Pezzella, L. Guarino, A. Piscitelli, How to enjoy laccases, *Cell Mol. Life Sci.* 72 (5) (2015) 923–940.
- [19] J. Chapman, A. Ismail, C. Dinu, Industrial applications of enzymes: recent advances, techniques, and outlooks, *Catalysts* 8 (2018) 238.
- [20] T.E. Sintra, S.P.M. Ventura, J.A.P. Coutinho, Superactivity induced by micellar systems as the key for boosting the yield of enzymatic reactions, *J. Mol. Catal. B Enzym.* 107 (2014) 140–151.
- [21] P.C.A.G. Pinto, M.L.M.F.S. Saraiva, J.L.F.C. Lima, Oxidoreductase behavior in ionic liquids: a review, *Anal. Sci.* 24 (10) (2008) 1231–1238.
- [22] E. Cunha, M.L.C. Passos, P.C.A.G. Pinto, M.L.M.F.S. Saraiva, Improved activity of α -chymotrypsin in mixed micelles of cetyltrimethylammonium bromide (CTAB) and ionic liquids: a kinetic study resorting to sequential injection analysis, *Colloids Surf. B Biointerf.* 118 (2014) 172–178.
- [23] P. Scovazzo, A.E. Visser, J.H. Davis, R.D. Rogers, C.A. Koval, D.L. DuBois, et al., Supported ionic liquid membranes and facilitated ionic liquid membranes, in: *Ionic Liquids*, American Chemical Society, ACS Symposium Series, vol. 818, 2002, pp. 6–69.
- [24] T. Welton, Ionic liquids in catalysis, *Coord. Chem. Rev.* 248 (21) (2004) 2459–2477.
- [25] H. Zhao, Methods for stabilizing and activating enzymes in ionic liquids—a review, *J. Chem. Technol. Biot.* 85 (2010) 891–907.
- [26] U. Preiss, C. Jungnickel, J. Thöming, I. Krossing, J. Luczak, M. Diedenhofen, A. Klant, Predicting the critical micelle concentrations of aqueous solutions of ionic liquids and other ionic surfactants, *Chem. – Eur. J.* 15 (35) (2009) 8880–8885.
- [27] M. Goldfeder, A. Fishman, Modulating enzyme activity using ionic liquids or surfactants, *Appl. Microbiol. Biotechnol.* 98 (2) (2014) 545–554.
- [28] T.L. Greaves, C.J. Drummond, Ionic liquids as amphiphile self-assembly media, *Chem. Soc. Rev.* 37 (8) (2008) 1709–1726.
- [29] J. Luczak, J. Hupka, J. Thöming, C. Jungnickel, Self-organization of imidazolium ionic liquids in aqueous solution, *Colloids Surf. Physicochem. Eng. Asp.* 329 (3) (2008) 125–133.
- [30] M.V. Quental, M. Caban, M.M. Pereira, P. Stepnowski, J.A.P. Coutinho, M.G. Freire, Enhanced extraction of proteins using cholinium-based ionic liquids as phase-forming components of aqueous biphasic systems, *Biotechnol. J.* 10 (9) (2015).
- [31] U. More, Z. Vaid, P. Bhamoria, A. Kumar NIM, Effect of [Cnmim][Br] based ionic liquids on the aggregation behavior of tetradecyltrimethylammonium bromide in aqueous medium, *J. Solut. Chem.* 44 (2015) 850–874.
- [32] E.L.P. de Faria, M.V. Gomes, A.F.M. Claudio, C.S.R. Freire, A.J.D. Silvestre, M.G. Freire, Extraction and recovery processes for cynaropicrin from *Cynara cardunculus* L. using aqueous solutions of surface-active ionic liquids, *Biophys. Rev.* 10 (3) (2018) 915–925.
- [33] F.A. Vicente, I.S. Cardoso, T.E. Sintra, J. Lemus, E.F. Marques, S.P.M. Ventura, et al., Impact of surface active ionic liquids on the cloud points of nonionic surfactants and the formation of aqueous micellar two-phase systems, *J. Phys. Chem. B* 121 (37) (2017) 8742–8755.
- [34] S. Kim, P.A. Thiessen, E.E. Bolton, J. Chen, G. Fu, A. Gindulyte, L. Han, J. He, S. He, B.A. Shoemaker, J. Wang, B. Yu, J. Zhang SHB, PubChem Substance and Compound databases.
- [35] O. Rodriguez, R.O. Cristovao, A.P.M. Tavares, E.A. Macedo, Study of the alkyl chain length on laccase stability and enzymatic kinetic with imidazolium ionic liquids, *Appl. Biochem. Biotechnol.* 164 (4) (2011) 524–533.
- [36] F. van Rantwijk, R.A. Sheldon, *Biocatalysis in Ionic Liquids*, 2007 (cited 2018 Sep 25).
- [37] G. Savelli, N. Spreti, P. Di Profio, Enzyme activity and stability control by amphiphilic self-organizing systems in aqueous solutions, *Curr. Opin. Colloid Interface Sci.* 5 (1) (2000) 111–117.
- [38] H. Zhao, Protein stabilization and enzyme activation in ionic liquids: specific ion effects: protein stabilization and enzyme activation in ionic liquids, *J. Chem. Technol. Biotechnol.* 91 (1) (2016) 25–50.
- [39] H. Liu, X. Wu, J. Sun, S. Chen, Stimulation of laccase biocatalysis in ionic liquids: a review on recent progress, *Curr. Protein Pept. Sci.* 19 (1) (2017) 100–111.
- [40] Z. Yang, Hofmeister effects: an explanation for the impact of ionic liquids on biocatalysis, *J. Chem. Technol. Biotechnol.* 144 (1) (2009) 12–22.
- [41] R. Campos, A. Kandelbauer, K.H. Robra, A. Cavaco-Paulo, G.M. Gübitz, Indigo degradation with purified laccases from *Trametes hirsuta* and *Sclerotium rolfsii*, *J. Biotechnol.* 89 (2) (2001) 131–139.
- [42] C.F. Thurston, The structure and function of fungal laccases, *Microbiology* 140 (1994) 19–26.
- [43] E. Ortiz, V. Gómez-Chávez, C.M. Cortés-Romero, H. Solís, S.L.-S. Rubén Ruiz-Ramos, Degradation of indigo carmine using advanced oxidation processes: synergy effects and toxicological study, *J. Environ. Prot.* 7 (2016) 1693–1706.
- [44] K.-K.M. Ho, Degradation of indigo and indigo carmine with an enzyme isolated from bacteria strain ATCC. p. US5457043A.
- [45] S. Ben Younes, S. Sayadi, Detoxification of Indigo carmine using a combined treatment via a novel trimeric thermostable laccase and microbial consortium, *J. Mol. Catal. B Enzym.* 87 (2013) 62–68.
- [46] E. Abadulla, T. Tzanov, S. Costa, K.-H. Robra, A. Cavaco-Paulo, G.M. Gübitz, Decolorization and detoxification of textile dyes with a Laccase from *Trametes hirsuta*, *Appl. Environ. Microbiol.* 66 (2000) 3357.
- [47] P.-P. Champagne, J.A. Ramsay, Dye decolorization and detoxification by laccase immobilized on porous glass beads, *Bioresour. Technol.* 101 (7) (2010) 2230–2235.
- [48] A. Kunamneni, I. Ghazi, S. Camarero, A. Ballesteros, F.J. Plou, M. Alcalde, Decolorization of synthetic dyes by laccase immobilized on epoxy-activated carriers, *Process Biochem.* 43 (2) (2008) 169–178.
- [49] Y. Wong, J. Yu, Laccase-catalyzed decolorization of synthetic dyes, *Water Res.* 33 (16) (1999) 3512–3520.
- [50] H. Forootanfar, A. Moezzi, M. Aghaie-Khozani, Y. Mahmoudjanlou, A. Ameri, F. Niknejad, et al., Synthetic dye decolorization by three sources of fungal laccase, *Iran. J. Environ. Health Sci. Eng.* 9 (1) (2012) 27.
- [51] S. Shipovskov, H.Q.N. Gunaratne, K.R. Seddon, G. Stephens, Catalytic activity of laccases in aqueous solutions of ionic liquids, *Green Chem.* 10 (7) (2008) 806–810.
- [52] L. Rehmann, E. Ivanova, J.L. Ferguson, H.Q.N. Gunaratne, K.R. Seddon, G.M. Stephens, Measuring the effect of ionic liquids on laccase activity using a simple, parallel method, *Green Chem.* 14 (3) (2012) 725–733.
- [53] O.V. Kushch, O.I. Hordieieva, O.O. Zosenko, A.N. Shendrik, Comparison of N-hydroxy compounds as mediators in laccase-catalysed decolorization of indigo carmine, *ChemistrySelect.* 4 (13) (2019) 3905–3913.
- [54] G. Xu, J. Wang, Q. Yin, W. Fang, Y. Xiao, Z. Fang, Expression of a thermo- and alkali-philic fungal laccase in *Pichia pastoris* and its application, *Protein Expr. Purif.* 154 (2019) 16–24.
- [55] V. Gupta, S. Balda, N. Gupta, N. Capalash, P. Sharma, Functional substitution of domain 3 (T1 copper center) of a novel laccase with Cu ions, *Int. J. Biol. Macromol.* 123 (2019) 1052–1061.
- [56] C. Louis-Jeune, M.A. Andrade-Navarro, C. Perez-Iratxeta, Prediction of protein secondary structure from circular dichroism using theoretically derived spectra, *Proteins Struct. Funct. Bioinforma.* 80 (2) (2012) 374–381.
- [57] P. Bhamoria, K.S. Rao, T.J. Trivedi, A. Kumar, Biomimetic ionic liquid induced folding alterations in the structure of bovine serum albumin in aqueous medium, *J. Phys. Chem. B* 118 (1) (2014) 115–124.
- [58] P. Bhamoria, M.J. Mehta, I. Pancha, A. Kumar, Structural and functional stability of cellulase in aqueous-biomimetic ionic liquid surfactant solution, *J. Phys. Chem. B* 118 (33) (2014) 9890–9899.
- [59] P. Bhamoria, T.J. Trivedi, A. Pabbathi, A. Samanta, A. Kumar, Ionic liquid-induced all- α to α + β conformational transition in cytochrome c with improved peroxidase activity in aqueous medium, *Phys. Chem. Chem. Phys.* 17 (15) (2015) 10189–10199.
- [60] P. Bhamoria, A. Kumar, Unusually high thermal stability and peroxidase activity of cytochrome c in ionic liquid colloidal formulation, *Chem. Commun.* 52 (3) (2016) 497–500.
- [61] Y. Liang, X. Yuan, G. Zeng, H. Zhong, H. Li W.W, Effects of surfactants on enzyme-containing reversed micellar system, *Sci. China Chem.* 54 (2011) 715.
- [62] R. Bianchini, G. Cevasco, C. Chiappe, C.S. Pomelli, M.J. Rodríguez Douton, Ionic liquids can significantly improve textile dyeing: an innovative application assuring economic and environmental benefits, *ACS Sustain. Chem. Eng.* 3 (9) (2015) 2303–2308.